

**Reference list for published genome-wide linkage scans  
for body weight and body fat in mice.**

- Anunciado, R.V., M. Nishimura, M. Mori, A. Ishikawa, S. Tanaka, F. Horio, T. Ohno, and T. Namikawa. 2001. Quantitative trait loci for body weight in the intercross between SM/J and A/J mice. *Exp Anim* **50**: 319-324.
- Brockmann, G.A., C.S. Haley, U. Renne, S.A. Knott, and M. Schwerin. 1998. Quantitative trait loci affecting body weight and fatness from a mouse line selected for extreme high growth. *Genetics* **150**: 369-381.
- Brockmann, G.A., E. Karatayli, C.S. Haley, U. Renne, O.J. Rottmann, and S. Karle. 2004. QTLs for pre- and postweaning body weight and body composition in selected mice. *Mamm Genome* **15**: 593-609.
- Brockmann, G.A., J. Kratzsch, C.S. Haley, U. Renne, M. Schwerin, and S. Karle. 2000. Single QTL effects, epistasis, and pleiotropy account for two-thirds of the phenotypic F(2) variance of growth and obesity in DU6i x DBA/2 mice. *Genome Res* **10**: 1941-1957.
- Cheverud, J.M., E.J. Routman, F.A. Duarte, B. van Swinderen, K. Cothran, and C. Perel. 1996. Quantitative trait loci for murine growth. *Genetics* **142**: 1305-1319.
- Cheverud, J.M., T.T. Vaughn, L.S. Pletscher, A.C. Peripato, E.S. Adams, C.F. Erikson, and K.J. King-Ellison. 2001. Genetic architecture of adiposity in the cross of LG/J and SM/J inbred mice. *Mamm Genome* **12**: 3-12.
- Corva, P.M., S. Horvat, and J.F. Medrano. 2001. Quantitative trait loci affecting growth in high growth (hg) mice. *Mamm Genome* **12**: 284-290.
- Dragani, T.A., Z.B. Zeng, F. Canzian, M. Gariboldi, M.T. Ghilarducci, G. Manenti, and M.A. Pierotti. 1995. Mapping of body weight loci on mouse chromosome X. *Mamm Genome* **6**: 778-781.
- Elliott, R.W., D. Poslinski, D. Tabaczynski, C. Hohman, and J. Pazik. 2004. Loci affecting male fertility in hybrids between *Mus macedonicus* and C57BL/6. *Mamm Genome* **15**: 704-710.
- Hirayama, I., Z. Yi, S. Izumi, I. Arai, W. Suzuki, Y. Nagamachi, H. Kuwano, T. Takeuchi, and T. Izumi. 1999. Genetic analysis of obese diabetes in the TSOD mouse. *Diabetes* **48**: 1183-1191.
- Horvat, S., L. Bunger, V.M. Falconer, P. Mackay, A. Law, G. Bulfield, and P.D. Keightley. 2000. Mapping of obesity QTLs in a cross between mouse lines divergently selected on fat content. *Mamm Genome* **11**: 2-7.
- Ishikawa, A., Y. Matsuda, and T. Namikawa. 2000. Detection of quantitative trait loci for body weight at 10 weeks from Philippine wild mice. *Mamm Genome* **11**: 824-830.
- Ishikawa, A. and T. Namikawa. 2004. Mapping major quantitative trait loci for postnatal growth in an intersubspecific backcross between C57BL/6J and Philippine wild mice by using principal component analysis. *Genes Genet Syst* **79**: 27-39.
- Ishimori, N., R. Li, P.M. Kelmenson, R. Korstanje, K.A. Walsh, G.A. Churchill, K. Forsman-Semb, and B. Paigen. 2004. Quantitative trait loci that determine plasma lipids and obesity in C57BL/6J and 129S1/SvImJ inbred mice. *J Lipid Res* **45**: 1624-1632.

- Keightley, P.D., T. Hardge, L. May, and G. Bulfield. 1996. A genetic map of quantitative trait loci for body weight in the mouse. *Genetics* **142**: 227-235.
- Keightley, P.D., K.H. Morris, A. Ishikawa, V.M. Falconer, and F. Oliver. 1998. Test of candidate gene--quantitative trait locus association applied to fatness in mice. *Heredity* **81 ( Pt 6)**: 630-637.
- Kluge, R., K. Giesen, G. Bahrenberg, L. Plum, J.R. Ortlepp, and H.G. Joost. 2000. Quantitative trait loci for obesity and insulin resistance (Nob1, Nob2) and their interaction with the leptin receptor allele (LeprA720T/T1044I) in New Zealand obese mice. *Diabetologia* **43**: 1565-1572.
- Lembertas, A.V., L. Perusse, Y.C. Chagnon, J.S. Fisler, C.H. Warden, D.A. Purcell-Huynh, F.T. Dionne, J. Gagnon, A. Nadeau, A.J. Lusis et al. 1997. Identification of an obesity quantitative trait locus on mouse chromosome 2 and evidence of linkage to body fat and insulin on the human homologous region 20q. *J Clin Invest* **100**: 1240-1247.
- Mehrabian, M., P.Z. Wen, J. Fisler, R.C. Davis, and A.J. Lusis. 1998. Genetic loci controlling body fat, lipoprotein metabolism, and insulin levels in a multifactorial mouse model. *J Clin Invest* **101**: 2485-2496.
- Moody, D.E., D. Pomp, M.K. Nielsen, and L.D. Van Vleck. 1999. Identification of quantitative trait loci influencing traits related to energy balance in selection and inbred lines of mice. *Genetics* **152**: 699-711.
- Morris, K.H., A. Ishikawa, and P.D. Keightley. 1999. Quantitative trait loci for growth traits in C57BL/6J x DBA/2J mice. *Mamm Genome* **10**: 225-228.
- Reed, D.R., X. Li, A.H. McDaniel, K. Lu, S. Li, M.G. Tordoff, R.A. Price, and A.A. Bachmanov. 2003. Loci on chromosomes 2, 4, 9, and 16 for body weight, body length, and adiposity identified in a genome scan of an F2 intercross between the 129P3/J and C57BL/6ByJ mouse strains. *Mamm Genome* **14**: 302-313.
- Reifsnyder, P.C., G. Churchill, and E.H. Leiter. 2000. Maternal environment and genotype interact to establish diabetes in mice. *Genome Res* **10**: 1568-1578.
- Rocha, J.L., E.J. Eisen, L.D. Van Vleck, and D. Pomp. 2004. A large-sample QTL study in mice: II. Body composition. *Mamm Genome* **15**: 100-113.
- Smith Richards, B.K., B.N. Belton, A.C. Poole, J.J. Mancuso, G.A. Churchill, R. Li, J. Volaufova, A. Zuberi, and B. York. 2002. QTL analysis of self-selected macronutrient diet intake: fat, carbohydrate, and total kilocalories. *Physiol Genomics* **11**: 205-217.
- Stoehr, J.P., J.E. Byers, S.M. Clee, H. Lan, I.V. Boronenkov, K.L. Schueler, B.S. Yandell, and A.D. Attie. 2004. Identification of major quantitative trait loci controlling body weight variation in ob/ob mice. *Diabetes* **53**: 245-249.
- Suto, J., S. Matsuura, K. Imamura, H. Yamanaka, and K. Sekikawa. 1998. Genetics of obesity in KK mouse and effects of A(y) allele on quantitative regulation. *Mamm Genome* **9**: 506-510.
- Taylor, B.A. and S.J. Phillips. 1996. Detection of obesity QTLs on mouse chromosomes 1 and 7 by selective DNA pooling. *Genomics* **34**: 389-398.
- Taylor, B.A. and S.J. Phillips. 1997. Obesity QTLs on mouse chromosomes 2 and 17. *Genomics* **43**: 249-257.
- Taylor, B.A., L.M. Tarantino, and S.J. Phillips. 1999. Gender-influenced obesity QTLs identified in a cross involving the KK type II diabetes-prone mouse strain. *Mamm Genome* **10**: 963-968.

- Taylor, B.A., C. Wnek, D. Schroeder, and S.J. Phillips. 2001. Multiple obesity QTLs identified in an intercross between the NZO (New Zealand obese) and the SM (small) mouse strains. *Mamm Genome* **12**: 95-103.
- Warden, C.H., J.S. Fisler, M.J. Pace, K.L. Svenson, and A.J. Lusis. 1993. Coincidence of genetic loci for plasma cholesterol levels and obesity in a multifactorial mouse model. *J Clin Invest* **92**: 773-779.
- Warden, C.H., J.S. Fisler, S.M. Shoemaker, P.Z. Wen, K.L. Svenson, M.J. Pace, and A.J. Lusis. 1995. Identification of four chromosomal loci determining obesity in a multifactorial mouse model. *J Clin Invest* **95**: 1545-1552.
- West, D.B., J. Goudey-Lefevre, B. York, and G.E. Truett. 1994a. Dietary obesity linked to genetic loci on chromosomes 9 and 15 in a polygenic mouse model. *J Clin Invest* **94**: 1410-1416.
- West, D.B., J. Waguespack, B. York, J. Goudey-Lefevre, and R.A. Price. 1994b. Genetics of dietary obesity in AKR/J x SWR/J mice: segregation of the trait and identification of a linked locus on chromosome 4. *Mamm Genome* **5**: 546-552.
- York, B., K. Lei, and D.B. West. 1997. Inherited non-autosomal effects on body fat in F2 mice derived from an AKR/J x SWR/J cross. *Mamm Genome* **8**: 726-730.
- Zhang, S. and H.K. Gershenfeld. 2003. Genetic contributions to body weight in mice: relationship of exploratory behavior to weight. *Obes Res* **11**: 828-838.